

Colorado Estimates OF People With TBI by Planning Region

CB Eagye, MS

Gale Whiteneck, PhD, FACRM

Craig Hospital

November 4, 2016

COLORADO PLANNING AND MANAGEMENT REGIONS




Table of Contents

- Planning Region 1 Page 26
- Planning Region 2 Page 53
- Planning Region 3 Page 80
- Planning Region 4 Page 107
- Planning Region 5 Page 134
- Planning Region 6 Page 161
- Planning Region 7 Page 188
- Planning Region 8 Page 215
- Planning Region 9 Page 242
- Planning Region 10 Page 269
- Planning Region 11 Page 296
- Planning Region 12 Page 323
- Planning Region 13 Page 350
- Planning Region 14 Page 377

Background

Current research built on the Colorado Study of TBI Prevalence and Associated Outcomes which was:


- CDC funded
- Statewide population-based survey of a random member of random-digit-dialed households.
- 2,701 adults age 18 and over interviewed using a modified CATI version of the OSU-TBI-ID
- To gather the lifetime history of TBI
- And standardized current assessments of 11 post-concussive symptoms and 10 negative outcomes


Raking and Weighting

- Raked the characteristics of the 2,701 people interviewed to the characteristics of the Colorado population age 18 and over
- Based on 2010 Census characteristics (including age; gender; race/ethnicity; marital, education, and employment status; and household income)
- Starting with a base weight of 1.0
- And iterating until all characteristics matched

Colorado Study Publications

- Whiteneck GG, Cuthbert JP, Corrigan JD, Bogner JA. Prevalence of Self-Reported Lifetime History of Traumatic Brain Injury and Associated Disability: A Statewide Population-Based Survey. J Head Trauma Rehabil. 2016 Jan-Feb;31(1):E55-62.
- Whiteneck GG, Cuthbert JP, Corrigan JD, Bogner JA. Risk of Negative Outcomes After Traumatic Brain Injury: A Statewide Population-Based Survey. J Head Trauma Rehabil. 2016 Jan-Feb;31(1):E43-54.
- Cuthbert JP, Whiteneck GG, Corrigan JD, Bogner J. The Reliability of a Computer-Assisted Telephone Interview Version of the Ohio State University Traumatic Brain Injury Identification Method. J Head Trauma Rehabil. 2016 Jan-Feb;31(1):E36-42.






Current Research Objectives

- To estimate the number of adults with a lifetime history of TBI
- And estimates of the numbers of people with TBI and 10 current negative outcomes as well as 11 post-concussive symptoms
- In each of 14 planning regions of Colorado
- To facilitate service delivery planning.

Current Research Methodology


- Secondary data analysis of the original study data
- Compare the population characteristics and rates of TBI in 3 large geographic areas of Colorado
 - The urban front-range counties
 - The western slope counties
 - The remaining non-urban, non-western slope counties in the east and south
- Use the respondents in these 3 large geographic areas to estimate TBI and outcomes in the 14 planning regions

Three Large Geographic Areas of Colorado




Statewide Data


Prevalence of TBI in Colorado




Statewide Injuries by Sex




Statewide Injuries by Age Groups




Statewide Injuries by Race




Gender Differences in Colorado



Age Differences in Colorado




TBI Prevalence Differences in Colorado




Current Research Raking and Weighting

- Raked the characteristics of people interviewed in the 3 large areas to the characteristics of the population age 18 and over in each of the 14 planning districts
- Based on 2016 population age and gender distributions of each planning district provided by the Colorado Office of Demography
- With the base weight as the original study weight
- And iterating until the characteristics matched


Gender Differences in Planning Regions



Age Differences in Planning Regions



TBI Prevalence Differences in Planning Regions



Closing Cautions

- These are only estimates
- Based on the self-reports
- Of 2,701 adult Coloradans
- With no estimates about TBI in children
- And no ability to infer causation, just association
- But hopefully helpful to TBI advocates and TBI service providers

Thank You

gale@craighospital.org
peagye@craighospital.org



CRAIG

UNYIELDING DETERMINATION.
EMPOWERING LIVES.


3425 S. Clarkson Street, Englewood, CO 80113

craighospital.org


Planning Region 1

Planning Region 1 Age 18+: 6 Injury Groups


Total Population: 55,303




Planning Region 1 Population by Age




Planning Region 1 Population by Sex



Planning Region 1


Poor Outcomes Among People with TBI

Planning Region 1

Postconcussive Symptoms Among People with TBI


Planning Region 1

- Endorsement of any cognitive complaints on the Alertness Behavior Subscale of the Sickness Impact Profile




Planning Region 1

- Two or more days of poor mental health reported




Planning Region 1

- Two or more days of poor physical health reported




Planning Region 1

- Limited in any activities or use special equipment




Planning Region 1

- More than 2 drinks per day for men (1 for women) or 5+ drinks in one sitting




Planning Region 1

- Limited in the kind or amount of work you can do




Planning Region 1

- Below average score on Diener Satisfaction with Life Scale




Planning Region 1

■ General health is fair or poor




Planning Region 1

- Need help from another person with activities of daily living




Planning Region 1

- Need to be reminded or supervised when doing activities of daily living




Planning Region 1

- Reported tiredness several days or more per month




Planning Region 1

- Reported sleep disturbance several days or more per month




Planning Region 1

- Reported irritability several days or more per month




Planning Region 1

- Reported headaches several days or more per month




Planning Region 1

- Reported poor concentration several days or more per month




Planning Region 1

- Reported poor memory several days or more per month




Planning Region 1

- Reported noise/light sensitivity several days or more per month




Planning Region 1

■ Reported dizziness several days or more per month




Planning Region 1

- Reported tinnitus several days or more per month




Planning Region 1

- Reported tiredness several days or more per month



Planning Region 1


- Reported tiredness several days or more per month




Planning Region 2

Planning Region 2 Age 18+: 6 Injury Groups


Total Population: 483,070




Planning Region 2 Population by Age




Planning Region 2 Population by Sex



Planning Region 2


Poor Outcomes Among People with TBI

Planning Region 2

Postconcussive Symptoms Among People with TBI


Planning Region 2

- Alertness Behavior Subscale of the Sickness Impact Profile – endorsement of any of 16 items like “I have difficulty reasoning and solving problems.”




Planning Region 2

- Two or more days of poor mental health reported




Planning Region 2

- Two or more days of poor physical health reported




Planning Region 2

- Limited in any activities or use special equipment




Planning Region 2

- More than 2 drinks per day for men (1 for women) or 5+ drinks in one sitting




Planning Region 2

- Limited in the kind or amount of work you can do




Planning Region 2

- Below average score on Diener Satisfaction with Life Scale




Planning Region 2

■ General health is fair or poor




Planning Region 2

- Need help from another person with activities of daily living




Planning Region 2

- Need to be reminded or supervised when doing activities of daily living




Planning Region 2

■ Reported tiredness several days or more per month




Planning Region 2

- Reported sleep disturbance several days or more per month




Planning Region 2

- Reported irritability several days or more per month




Planning Region 2

- Reported headaches several days or more per month




Planning Region 2

- Reported poor concentration several days or more per month




Planning Region 2

- Reported poor memory several days or more per month




Planning Region 2

- Reported noise/light sensitivity several days or more per month




Planning Region 2

- Reported tiredness several days or more per month




Planning Region 2

- Reported tinnitus several days or more per month




Planning Region 2

- Reported poor balance several days or more per month



Planning Region 2


- Reported poor vision several days or more per month




Planning Region 3

Planning Region 3 Age 18+: 6 Injury Groups


Total Population: 2,397,584




Planning Region 3 Population by Age




Planning Region 3 Population by Sex



Planning Region 3


Poor Outcomes Among People with TBI

Planning Region 3

Postconcussive Symptoms Among People with TBI


Planning Region 3

- Endorsement of any cognitive complaints on the Alertness Behavior Subscale of the Sickness Impact Profile




Planning Region 3

- Two or more days of poor mental health reported




Planning Region 3

■ Two or more days of poor physical health reported




Planning Region 3

■ Limited in any activities or use special equipment




Planning Region 3

- More than 2 drinks per day for men (1 for women) or 5+ drinks in one sitting




Planning Region 3

- Limited in the kind or amount of work you can do




Planning Region 3

- Below average score on Diener Satisfaction with Life Scale




Planning Region 3

■ General health is fair or poor




Planning Region 3

- Need help from another person with activities of daily living




Planning Region 3

- Need to be reminded or supervised when doing activities of daily living




Planning Region 3

- Reported tiredness several days or more per month




Planning Region 3

■ Reported sleep disturbance several days or more per month




Planning Region 3

- Reported irritability several days or more per month




Planning Region 3

- Reported headaches several days or more per month




Planning Region 3

■ Reported poor concentration several days or more per month




Planning Region 3

- Reported poor memory several days or more per month




Planning Region 3

■ Reported noise/light sensitivity several days or more per month




Planning Region 3

■ Reported dizziness several days or more per month




Planning Region 3

- Reported tinnitus several days or more per month




Planning Region 3

- Reported poor balance several days or more per month



Planning Region 3


■ Reported poor vision several days or more per month




Planning Region 4

Planning Region 4 Age 18+: 6 Injury Groups


Total Population: 544,965




Planning Region 4 Population by Age




Planning Region 4 Population by Sex



Planning Region 4


Poor Outcomes Among People with TBI

Planning Region 4

Postconcussive Symptoms Among People with TBI


Planning Region 4

- Endorsement of any cognitive complaints on the Alertness Behavior Subscale of the Sickness Impact Profile




Planning Region 4

- Two or more days of poor mental health reported




Planning Region 4

- Two or more days of poor physical health reported




Planning Region 4

- Limited in any activities or use special equipment




Planning Region 4

- More than 2 drinks per day for men (1 for women) or 5+ drinks in one sitting




Planning Region 4

- Limited in the kind or amount of work you can do




Planning Region 4

- Below average score on Diener Satisfaction with Life Scale




Planning Region 4

■ General health is fair or poor




Planning Region 4

- Need help from another person with activities of daily living




Planning Region 4

- Need to be reminded or supervised when doing activities of daily living




Planning Region 4

■ Reported tiredness several days or more per month




Planning Region 4

- Reported sleep disturbance several days or more per month




Planning Region 4

■ Reported irritability several days or more per month




Planning Region 4

- Reported headaches several days or more per month




Planning Region 4

■ Reported poor concentration several days or more per month




Planning Region 4

■ Reported poor memory several days or more per month




Planning Region 4

- Reported noise/light sensitivity several days or more per month




Planning Region 4

■ Reported dizziness several days or more per month




Planning Region 4

■ Reported tinnitus several days or more per month




Planning Region 4

■ Reported poor balance several days or more per month



Planning Region 4


■ Reported poor vision several days or more per month




Planning Region 5

Planning Region 5 Age 18+: 6 Injury Groups


Total Population: 33,239




Planning Region 5 Population by Age




Planning Region 5 Population by Sex



Planning Region 5


Postconcussive Symptoms Among People with TBI

Planning Region 5

Poor Outcomes Among People with TBI


Planning Region 5

- Endorsement of any cognitive complaints on the Alertness Behavior Subscale of the Sickness Impact Profile




Planning Region 5

- Two or more days of poor mental health reported




Planning Region 5

- Two or more days of poor physical health reported




Planning Region 5

- Limited in any activities or use special equipment




Planning Region 5

- More than 2 drinks per day for men (1 for women) or 5+ drinks in one sitting




Planning Region 5

- Limited in the kind or amount of work you can do




Planning Region 5

■ Below average score on Diener Satisfaction with Life Scale




Planning Region 5

■ General health is fair or poor




Planning Region 5

■ Need help from another person with activities of daily living




Planning Region 5

- Need to be reminded or supervised when doing activities of daily living




Planning Region 5

- Reported tiredness several days or more per month




Planning Region 5

■ Reported sleep disturbance several days or more per month




Planning Region 5

■ Reported irritability several days or more per month




Planning Region 5

■ Reported headaches several days or more per month




Planning Region 5

- Reported poor concentration several days or more per month




Planning Region 5

■ Reported poor memory several days or more per month




Planning Region 5

- Reported noise/light sensitivity several days or more per month




Planning Region 5

- Reported dizziness several days or more per month




Planning Region 5

■ Reported tinnitus several days or more per month




Planning Region 5

- Reported poor balance several days or more per month



Planning Region 5


■ Reported poor vision several days or more per month




Planning Region 6

Planning Region 6 Age 18+: 6 Injury Groups


Total Population: 36,674




Planning Region 6 Population by Age




Planning Region 6 Population by Sex



Planning Region 6


Poor Outcomes Among People with TBI

Planning Region 6

Postconcussive Symptoms Among People with TBI


Planning Region 6

- Endorsement of any cognitive complaints on the Alertness Behavior Subscale of the Sickness Impact Profile




Planning Region 6

- Two or more days of poor mental health reported




Planning Region 6

■ Two or more days of poor physical health reported




Planning Region 6

■ Limited in any activities or use special equipment




Planning Region 6

- More than 2 drinks per day for men (1 for women) or 5+ drinks in one sitting




Planning Region 6

- Limited in the kind or amount of work you can do




Planning Region 6

- Below average score on Diener Satisfaction with Life Scale




Planning Region 6

■ General health is fair or poor




Planning Region 6

- Need help from another person with activities of daily living




Planning Region 6

- Need to be reminded or supervised when doing activities of daily living




Planning Region 6

■ Reported tiredness several days or more per month




Planning Region 6

■ Reported sleep disturbance several days or more per month




Planning Region 6

- Reported irritability several days or more per month




Planning Region 6

■ Reported headaches several days or more per month




Planning Region 6

■ Reported poor concentration several days or more per month




Planning Region 6

■ Reported poor memory several days or more per month




Planning Region 6

- Reported noise/light sensitivity several days or more per month




Planning Region 6

- Reported dizziness several days or more per month




Planning Region 6

■ Reported tinnitus several days or more per month




Planning Region 6

■ Reported poor balance several days or more per month



Planning Region 6


■ Reported poor vision several days or more per month




Planning Region 7

Planning Region 7 Age 18+: 6 Injury Groups


Total Population: 126,073




Planning Region 7 Population by Age




Planning Region 7 Population by Sex



Planning Region 7


Postconcussive Symptoms Among People with TBI

Planning Region 7

Poor Outcomes Among People with TBI


Planning Region 7

- Endorsement of any cognitive complaints on the Alertness Behavior Subscale of the Sickness Impact Profile




Planning Region 7

■ Two or more days of poor mental health reported




Planning Region 7

■ Two or more days of poor physical health reported




Planning Region 7

- Limited in any activities or use special equipment




Planning Region 7

- More than 2 drinks per day for men (1 for women) or 5+ drinks in one sitting




Planning Region 7

- Limited in the kind or amount of work you can do




Planning Region 7

■ Below average score on Diener Satisfaction with Life Scale




Planning Region 7

■ General health is fair or poor




Planning Region 7

■ Need help from another person with activities of daily living




Planning Region 7

- Need to be reminded or supervised when doing activities of daily living




Planning Region 7

- Reported tiredness several days or more per month




Planning Region 7

■ Reported sleep disturbances several days or more per month




Planning Region 7

- Reported irritability several days or more per month




Planning Region 7

■ Reported headaches several days or more per month




Planning Region 7

■ Reported poor concentration several days or more per month




Planning Region 7

- Reported poor memory several days or more per month




Planning Region 7

■ Reported noise/light sensitivity several days or more per month




Planning Region 7

■ Reported dizziness several days or more per month




Planning Region 7

- Reported tinnitus several days or more per month




Planning Region 7

■ Reported poor balance several days or more per month



Planning Region 7


■ Reported poor vision several days or more per month




Planning Region 8

Planning Region 8 Age 18+: 6 Injury Groups


Total Population: 35,125




Planning Region 8 Population by Age




Planning Region 8 Population by Sex



Planning Region 8


Poor Outcomes Among People with TBI

Planning Region 8

Postconcussive Symptoms Among People with TBI


Planning Region 8

- Endorsement of any cognitive complaints on the Alertness Behavior Subscale of the Sickness Impact Profile




Planning Region 8

- Two or more days of poor mental health reported




Planning Region 8

■ Two or more days of poor physical health reported




Planning Region 8

- Limited in any activities or use special equipment




Planning Region 8

- More than 2 drinks per day for men (1 for women) or 5+ drinks in one sitting




Planning Region 8

- Limited in the kind or amount of work you can do




Planning Region 8

- Below average score on Diener Satisfaction with Life Scale




Planning Region 8

■ General health is fair or poor




Planning Region 8

- Need help from another person with activities of daily living




Planning Region 8

- Need to be reminded or supervised when doing activities of daily living




Planning Region 8

- Reported tiredness several days or more per month




Planning Region 8

■ Reported sleep disturbance several days or more per month




Planning Region 8

■ Reported irritability several days or more per month




Planning Region 8

- Reported headaches several days or more per month




Planning Region 8

■ Reported poor concentration several days or more per month




Planning Region 8

■ Reported poor memory several days or more per month




Planning Region 8

- Reported noise/light sensitivity several days or more per month




Planning Region 8

■ Reported dizziness several days or more per month




Planning Region 8

■ Reported tinnitus several days or more per month




Planning Region 8

■ Reported poor balance several days or more per month



Planning Region 8


■ Reported poor vision several days or more per month




Planning Region 9

Planning Region 9 Age 18+: 6 Injury Groups


Total Population: 33,265




Planning Region 9 Population by Age




Planning Region 9 Population by Sex



Planning Region 9


Postconcussive Symptoms Among People with TBI

Planning Region 9

Poor Outcomes Among People with TBI


Planning Region 9

- Endorsement of any cognitive complaints on the Alertness Behavior Subscale of the Sickness Impact Profile




Planning Region 9

■ Two or more days of poor mental health reported




Planning Region 9

■ Two or more days of poor physical health reported




Planning Region 9

- Limited in any activities or use special equipment




Planning Region 9

- More than 2 drinks per day for men (1 for women) or 5+ drinks in one sitting




Planning Region 9

- Limited in the kind or amount of work you can do




Planning Region 9

■ Below average score on Diener Satisfaction with Life Scale




Planning Region 9

■ General health is fair or poor




Planning Region 9

- Need help from another person with activities of daily living




Planning Region 9

- Need to be reminded or supervised when doing activities of daily living




Planning Region 9

■ Reported tiredness several days or more per month




Planning Region 9

■ Reported sleep disturbance several days or more per month




Planning Region 9

■ Reported irritability several days or more per month




Planning Region 9

- Reported headaches several days or more per month




Planning Region 9

■ Reported poor concentration several days or more per month




Planning Region 9

- Reported poor memory several days or more per month




Planning Region 9

- Reported noise/light sensitivity several days or more per month




Planning Region 9

■ Reported dizziness several days or more per month




Planning Region 9

■ Reported tinnitus several days or more per month




Planning Region 9

- Reported poor balance several days or more per month



Planning Region 9


■ Reported poor vision several days or more per month



Planning Region 10


Planning Region 10 Age 18+: 6 Injury Groups

Total Population: 81,465




- No Injury - 16,154
- Injury No TBI - 31,335
- Mild TBI No LOC - 8,234
- Mild TBI with LOC - 20,062
- Moderate TBI - 4,376
- Severe TBI - 1,304

Planning Region 10 Population by Age




Planning Region 10 Population by Sex




Planning Region 10

Postconcussive Symptoms Among People with TBI




Planning Region 10

Poor Outcomes Among People with TBI


Planning Region 10

- Endorsement of any cognitive complaints on the Alertness Behavior Subscale of the Sickness Impact Profile




Planning Region 10

- Two or more days of poor mental health reported




Planning Region 10

- Two or more days of poor physical health reported




Planning Region 10

- Limited in any activities or use special equipment




Planning Region 10

- More than 2 drinks per day for men (1 for women) or 5+ drinks in one sitting




Planning Region 10

- Limited in the kind or amount of work you can do




Planning Region 10

■ Below average score on Diener Satisfaction with Life Scale




Planning Region 10

■ General health is fair or poor




Planning Region 10

- Need help from another person with activities of daily living




Planning Region 10

- Need to be reminded or supervised when doing activities of daily living




Planning Region 10

- Reported tiredness several days or more per month




Planning Region 10

■ Reported sleep disturbance several days or more per month




Planning Region 10

- Reported irritability several days or more per month




Planning Region 10

■ Reported headaches several days or more per month




Planning Region 10

■ Reported poor concentration several days or more per month




Planning Region 10

■ Reported poor memory several days or more per month




Planning Region 10

- Reported noise/light sensitivity several days or more per month




Planning Region 10

- Reported dizziness several days or more per month




Planning Region 10

■ Reported tinnitus several days or more per month




Planning Region 10

■ Reported poor balance several days or more per month



Planning Region 10


■ Reported poor vision several days or more per month




Planning Region 11

Planning Region 11 Age 18+: 6 Injury Groups


Total Population: 195,773




Planning Region 11 Population by Age




Planning Region 11 Population by Sex



Planning Region 11


Postconcussive Symptoms Among People with TBI

Planning Region 11

Poor Outcomes Among People with TBI


Planning Region 11

- Endorsement of any cognitive complaints on the Alertness Behavior Subscale of the Sickness Impact Profile




Planning Region 11

■ Two or more days of poor mental health reported




Planning Region 11

■ Two or more days of poor physical health reported




Planning Region 11

■ Limited in any activities or use special equipment




Planning Region 11

- More than 2 drinks per day for men (1 for women) or 5+ drinks in one sitting




Planning Region 11

- Limited in the kind or amount of work you can do




Planning Region 11

■ Below average score on Diener Satisfaction with Life Scale




Planning Region 11

■ General health is fair or poor




Planning Region 11

- Need help from another person with activities of daily living




Planning Region 11

- Need to be reminded or supervised when doing activities of daily living




Planning Region 11

■ Reported tiredness several days or more per month




Planning Region 11

■ Reported sleep disturbance several days or more per month




Planning Region 11

- Reported irritability several days or more per month




Planning Region 11

■ Reported headaches several days or more per month




Planning Region 11

■ Reported poor concentration several days or more per month




Planning Region 11

■ Reported poor memory several days or more per month




Planning Region 11

- Reported noise/light sensitivity several days or more per month




Planning Region 11

■ Reported dizziness several days or more per month




Planning Region 11

- Reported tinnitus several days or more per month




Planning Region 11

■ Reported poor balance several days or more per month



Planning Region 11


■ Reported poor vision several days or more per month




Planning Region 12

Planning Region 12 Age 18+: 6 Injury Groups


Total Population: 94,178




Planning Region 12 Population by Age




Planning Region 12 Population by Sex



Planning Region 12


Poor Outcomes Among People with TBI

Planning Region 12

Postconcussive Symptoms Among People with TBI


Planning Region 12

- Endorsement of any cognitive complaints on the Alertness Behavior Subscale of the Sickness Impact Profile




Planning Region 12

■ Two or more days of poor mental health reported




Planning Region 12

■ Two or more days of poor physical health reported




Planning Region 12

- Limited in any activities or use special equipment




Planning Region 12

- More than 2 drinks per day for men (1 for women) or 5+ drinks in one sitting




Planning Region 12

- Limited in the kind or amount of work you can do




Planning Region 12

- Below average score on Diener Satisfaction with Life Scale




Planning Region 12

■ General health is fair or poor




Planning Region 12

- Need help from another person with activities of daily living




Planning Region 12

- Need to be reminded or supervised when doing activities of daily living




Planning Region 12

■ Reported tiredness several days or more per month




Planning Region 12

■ Reported sleep disturbance several days or more per month




Planning Region 12

■ Reported irritability several days or more per month




Planning Region 12

■ Reported headaches several days or more per month




Planning Region 12

■ Reported poor concentration several days or more per month




Planning Region 12

■ Reported poor memory several days or more per month




Planning Region 12

- Reported noise/light sensitivity several days or more per month




Planning Region 12

■ Reported dizziness several days or more per month




Planning Region 12

■ Reported tinnitus several days or more per month




Planning Region 12

- Reported poor balance several days or more per month



Planning Region 12


■ Reported poor vision several days or more per month




Planning Region 13

Planning Region 13 Age 18+: 6 Injury Groups


Total Population: 110,017




Planning Region 13 Population by Age




Planning Region 13 Population by Sex



Planning Region 13


Postconcussive Symptoms Among People with TBI

Planning Region 13

Poor Outcomes Among People with TBI


Planning Region 13

■ Endorsement of any cognitive complaints on the Alertness Behavior Subscale of the Sickness Impact Profile




Planning Region 13

■ Two or more days of poor mental health reported




Planning Region 13

■ Two or more days of poor physical health reported




Planning Region 13

- Limited in any activities or use special equipment




Planning Region 13

- More than 2 drinks per day for men (1 for women) or 5+ drinks in one sitting




Planning Region 13

- Limited in the kind or amount of work you can do




Planning Region 13

- Below average score on Diener Satisfaction with Life Scale




Planning Region 13

■ General health is fair or poor




Planning Region 13

- Need help from another person with activities of daily living




Planning Region 13

- Need to be reminded or supervised when doing activities of daily living




Planning Region 13

■ Reported tiredness several days or more per month




Planning Region 13

■ Reported sleep disturbance several days or more per month




Planning Region 13

■ Reported tiredness several days or more per month




Planning Region 13

■ Reported headaches several days or more per month




Planning Region 13

■ Reported poor concentration several days or more per month




Planning Region 13

■ Reported poor memory several days or more per month




Planning Region 13

- Reported noise/light sensitivity several days or more per month




Planning Region 13

■ Reported dizziness several days or more per month




Planning Region 13

- Reported tinnitus several days or more per month




Planning Region 13

■ Reported poor balance several days or more per month



Planning Region 13


■ Reported poor vision several days or more per month




Planning Region 14

Planning Region 14 Age 18+: 6 Injury Groups


Total Population: 17,151




Planning Region 14 Population by Age




Planning Region 14 Population by Sex



Planning Region 14


Postconcussive Symptoms Among People with TBI

Planning Region 14

Poor Outcomes Among People with TBI


Planning Region 14

- Endorsement of any cognitive complaints on the Alertness Behavior Subscale of the Sickness Impact Profile




Planning Region 14

■ Two or more days of poor mental health reported




Planning Region 14

■ Two or more days of poor physical health reported




Planning Region 14

- Limited in any activities or use special equipment




Planning Region 14

- More than 2 drinks per day for men (1 for women) or 5+ drinks in one sitting




Planning Region 14

- Limited in the kind or amount of work you can do




Planning Region 14

- Below average score on Diener Satisfaction with Life Scale




Planning Region 14

■ General health is fair or poor




Planning Region 14

- Need help from another person with activities of daily living




Planning Region 14

- Need to be reminded or supervised when doing activities of daily living




Planning Region 14

■ Reported tiredness several days or more per month




Planning Region 14

- Reported sleep disturbance several days or more per month




Planning Region 14

■ Reported irritability several days or more per month




Planning Region 14

■ Reported headaches several days or more per month




Planning Region 14

■ Reported poor concentration several days or more per month




Planning Region 14

■ Reported poor memory several days or more per month




Planning Region 14

■ Reported noise/light sensitivity several days or more per month




Planning Region 14

■ Reported dizziness several days or more per month




Planning Region 14

- Reported tinnitus several days or more per month



Planning Region 14

- Reported poor balance several days or more per month



Planning Region 14

■ Reported poor vision several days or more per month

